

## Diurnal Birds (Animalia: Aves) in the Area of Taman Wisata Alam Gunung Meja, Manokwari, West Papua

### Burung Diurnal (Animalia: Aves) di Kawasan Taman Wisata Alam Gunung Meja, Manokwari, Papua Barat

Freddy Pattiselanno<sup>1\*</sup>, Johan F. Koibur<sup>1</sup>, Hotlan Manik<sup>1</sup>, and Agustina Y.S. Arobaya<sup>2</sup>

<sup>1</sup>Sub-Laboratorium Budidaya Aneka Ternak dan Satwa

Fakultas Peternakan Perikanan dan Ilmu Kelautan Universitas Negeri Papua, Manokwari

<sup>2</sup>Pusat Penelitian Keanekaragaman Hayati Universitas Negeri Papua, Manokwari

E-mail: freddy.pattiselanno@fppk.unipa.ac.id \* Corresponding author

Taman Wisata Alam Gunung Meja (TWGM) located in 134°03'17"–134°04'05" E and 0°51'29"–0°52'59" S, is one of unique protected areas in Papua, categorized as a lowland tropical rainforest represented holotype forest areas in the coastal site of the northern Papua.

Survey was conducted in the area of TWGM from 8 to 22 November 2006 by daily walking along 1,5km constructing transect trail fit to the *topography condition*. In general, all observation sites are grown by woody plants, palms, lianas, ferns, and shrubs species. The plants species around the sites were not observed in details, however, some woody species had been recognized such as *Pometia pinnata*, *Vitex confassus*, *Horfieldia irya*, *Sterculia macrophylla*, *Pimelodendron amboinicum*, *Palaquium amboinensis*, *Horsfieldia silvetris* and *Aglaia* sp. Others like *Bambusa* sp, palms (*Licuala* sp., *Calamus aruensis* Becc and *Caryota rumphiana* Mart); lianas (*Asplenium nidus* L., *Acrosticum aureum* L., *Selaginela australis* and *Dipteris steroides*); ferns (*Dicksonia blumei* Moore, *Lycopodium complanatum* L. and *Asplenidium nidus* L.; shrubs (*Mimosa pudica*, *Euphorbia geniculata*, *Bidens pilosa*, *Imperata cylindrica*, *Paspalum conjugata*, *Crotalaria incana* L. *Laucaena glauca* and *Axonopus compressus*); orchids (*Bulbohyllum* sp., *Spathoglotis* sp. And *Dendrobium littorale*) were also encountered during the survey. Topography was the only difference between two transect lines. Line 1 was done along the road built within the areas (the main road is usually used to pass the area from one to the other site). Therefore, this line

had a fair and light topography, and it was easy to conduct. Along the line, woody plants were grown with rare canopy cover, so the sun light could easily penetrate the low layer forest land. On the other hand, transect line 2 was established within the forest site, entering deeper, the study site was covered by woody and high trees, so most of the sites were fully covered with canopy. It had heavy topography with constricted relief. The site was more humid, and it was found that the temperature was lower (25°C in average) compared to transect 1 (28°C in average) while the forest floor was sometimes wet. Bird observation was performed twice a day, early in the morning between 05.00–07.00 a.m. and late in the afternoon between 04.00–06.00 p.m. An additional observation was also carried out in two different observation points (1) 15m asl 134°04'359"E and 00°50'753"S and (2) 17m asl 134°04'485"E and 00°50'981"S. These two observation points were differentiated from the situation of the sites; where point 1 was commonly surrounded by plantation areas (corns, cassava and some fruit plants), almost found at flat areas adjacent to local settlements and point 2 was secondary forests with some hilly parts. Birds were observed with the aid of Bushnell binocular 10x50, and identified using the taxonomic description from Rand and Gilliard (1967) and Beehler *et al.*, (2001). The number of species observed during the day survey was recorded and a species accumulation curve was constructed from daily totals. Species abundance was classified into five different categories: *Abundant* (recorded regularly, in moderate to large number);

*Common* (recorded regularly, in small number); *Fairly* (recorded scattered not as usual as common); *Uncommon* (recorded irregularly); and *Rare* (recorded once or twice) and tabulated in the species list (Table 1).

Forty five species belong to twenty two families were recorded during the survey, 32 species along the transect line, 22 species at point 1 and 27 species at point 2. Previous study on avifauna in the similar relative location, indicated that approximately 15 families represented 35 bird species were recorded, 14 were endemic species and 21 species non-endemic (Dwiyanto, 1995) and roughly 20 families composed of 35 species encountered during a series of observation published by the Forest Research Institute of Manokwari (Leppe and Tokede, 2006).

This study also revealed that common and abundant species, such as *C. tristis*, *D. pinon*, *G. chloronotus*, *H. indus*, *L. lorry*, *R. plicatus* and *R. rufiventris* were also found in the whole of New Guineas Island and its satellite island, like Raja Ampat (Mayor and Pattiselanno, 2008). While *E. ibis* for the last fifteen years were increased in Papua, a part of

its spread out distribution around the world was also found particularly in the warmer latitude sites (Beehler *et al.*, 2001). Particular species from certain families were also encountered by Pattiselanno (2005) during the survey carried out around the Mamberamo River Basin. *C. esculanta* was not only found in Papua, but also in whole Indonesia archipelago and Malaya. In contrast, some species were found rare, *S. spilodera*, *P. nanus*, *T. megalorynchos* and *R. leucothorax* that were only found in one of three observed sites.

A combined species accumulation curve for transect line and both two observation points (Figure 1) assumes that further observation efforts should be approached so significant results would be obtained. Various results gathered from previous surveys conducted to the relative similar location may be due to different methods and vocal point in observing the bird species. Therefore, using vocal recorded to support the data collection is important, because birds in tropical rainforest are difficult to observe. According to Beehler and Mack (1998), vocal repertoires and local dialects are vital in identifying bird.




Figure 1. Species accumulation curve for birds at sites around Taman Wisata Gunung Meja, Manokwari.

**Table 1.** Number of bird species observed along the transect and observation point.

| No. | Species (1)                    | Transect Line (2) | Point 1 (3) | Point 2 (4) | Species Abundance (5) | Remarks (6) |
|-----|--------------------------------|-------------------|-------------|-------------|-----------------------|-------------|
| 1.  | <b>Accipitridae</b>            |                   |             |             |                       |             |
|     | <i>Accipiter fasciatus</i>     | √                 | √           |             | Uncommon              |             |
|     | <i>A. novaehollandiae</i>      | √                 | √           | √           | Common                |             |
|     | <i>Circus spilonotus</i>       |                   |             | √           | Fairly                |             |
|     | <i>Haliastur indus</i>         | √                 | √           |             | Common                |             |
| 2.  | <b>Achantizidae</b>            |                   |             |             |                       |             |
|     | <i>Gerygone chloronotus</i>    | √                 | √           | √           | Abundant              | P           |
|     | <i>G. cinerea</i>              |                   | √           |             | Abundant              |             |
|     | <i>Crateroscelis nigrofura</i> | √                 |             | √           | Uncommon              |             |
|     | <i>Sericornis spilodera</i>    |                   |             | √           | Rare                  | P           |
| 3.  | <b>Alcedinidae</b>             |                   |             |             |                       |             |
|     | <i>Dacelo gaudichaud</i>       | √                 | √           | √           | Abundant              | E-NG; P     |
|     | <i>Halycon nigrocyanea</i>     | √                 |             | √           | Common                |             |
|     | <i>H. macleayi</i>             | √                 |             | √           | Abundant              | P           |
|     | <i>H. sancta</i>               | √                 | √           | √           | Abundant              |             |
| 4.  | <b>Apodidae</b>                |                   |             |             |                       |             |
|     | <i>Collocalia esculanta</i>    | √                 |             | √           | Abundant              |             |
| 5.  | <b>Ardeidae</b>                |                   |             |             |                       |             |
|     | <i>Egretta ibis</i>            | √                 |             | √           | Common                |             |
| 6.  | <b>Bucerotidae</b>             |                   |             |             |                       |             |
|     | <i>Rhyticeros plicatus</i>     | √                 | √           | √           | Abundant              |             |
| 7.  | <b>Campephagidae</b>           |                   |             |             |                       |             |
|     | <i>Coracina melaena</i>        |                   |             | √           | Uncommon              |             |
| 8.  | <b>Columbidae</b>              |                   |             |             |                       |             |
|     | <i>Ducula pinon</i>            | √                 | √           |             | Common                | E-NG        |
|     | <i>Ptilinopus nanus</i>        |                   |             | √           | Rare                  | E-NG        |
|     | <i>P. solomonensis</i>         | √                 |             |             | Uncommon              |             |
| 9.  | <b>Corvidae</b>                |                   |             |             |                       |             |
|     | <i>Corvus tristis</i>          | √                 |             | √           | Common                | E-NG        |
| 10. | <b>Cracticidae</b>             |                   |             |             |                       |             |
|     | <i>Cracticus quoyi</i>         | √                 | √           | √           | Abundant              |             |
| 11. | <b>Dicruridae</b>              |                   |             |             |                       |             |
|     | <i>Dicrurus huttentottus</i>   | √                 |             |             | Fairly                |             |
| 12. | <b>Estrildidae</b>             |                   |             |             |                       |             |
|     | <i>Lonchura tristissima</i>    | √                 | √           |             | Abundant              | E-NG        |
| 13. | <b>Falconidae</b>              |                   |             |             |                       |             |
|     | <i>Falco severus</i>           |                   |             | √           | Uncommon              |             |
| 14. | <b>Maluridae</b>               |                   |             |             |                       |             |
|     | <i>Malurus alboscapulatus</i>  | √                 | √           |             | Abundant              | E-NG        |
| 15. | <b>Meliphagidae</b>            |                   |             |             |                       |             |
|     | <i>Meliphaga montana</i>       |                   | √           | √           | Abundant              | E-NG        |
|     | <i>Oedistoma pygmaeum</i>      | √                 | √           | √           | Abundant              |             |
|     | <i>Philemon buceroides</i>     | √                 |             |             | Uncommon              |             |
|     | <i>Xanthotis flaviventer</i>   |                   |             | √           | Fairly                |             |
| 16. | <b>Myagridae</b>               |                   |             |             |                       |             |
|     | <i>Monarcha guttula</i>        | √                 |             | √           | Common                | E-NG        |
|     | <i>M. manadensis</i>           |                   | √           | √           | Common                | E-NG        |
| 17. | <b>Oriolidae</b>               |                   |             |             |                       |             |
|     | <i>Oriolus szalayii</i>        | √                 | √           |             | Common                | E-NG        |
| 18. | <b>Podargidae</b>              |                   |             |             |                       |             |
|     | <i>Podargus papuensis</i>      | √                 | √           | √           | Common                |             |

Diurnal birds (Animalia: Aves) at the area of Taman Wisata Alam Gunung Meja

Table 1. Continue

| No. | Species (1)                          | Transect Line (2) | Point 1 (3) | Point 2 (4) | Species Abundance (5) | Remarks (6) |
|-----|--------------------------------------|-------------------|-------------|-------------|-----------------------|-------------|
| 19. | <b>Psittacidae</b>                   |                   |             |             |                       |             |
|     | <i>Chamosyna placentis</i>           | √                 |             |             | Uncommon              |             |
|     | <i>Eclectus roratus</i>              | √                 | √           |             | Common                | P           |
|     | <i>Lorius lory</i>                   | √                 |             | √           | Abundant              | E-NG        |
|     | <i>Geoffroyus geoffroyi</i>          |                   | √           | √           | Abundant              |             |
|     | <i>Probosciger aterrimus</i>         | √                 |             |             | Uncommon              |             |
|     | <i>Psittaculirostris desmarestii</i> |                   | √           |             | Fairly                |             |
|     | <i>Tanygnathus megalorynchos</i>     | √                 |             |             | Rare                  |             |
| 20. | <b>Rhipiduridae</b>                  |                   |             |             |                       |             |
|     | <i>Rhipidura leucothorax</i>         | √                 |             |             | Rare                  | E-NG        |
|     | <i>R. rufiventris</i>                |                   | √           | √           | Common                |             |
| 21. | <b>Sturnidae</b>                     |                   |             |             |                       |             |
|     | <i>Mino dumontii</i>                 | √                 |             | √           | Uncommon              |             |
| 22. | <b>Zosteropidae</b>                  |                   |             |             |                       |             |
|     | <i>Zosterops fuscicapillus</i>       |                   | √           |             | Fairly                |             |
|     | <i>Z. novae guineae</i>              | √                 |             |             | Common                |             |

Note: √= present, E-NG= Endemic New Guinea, P= Protected

## Acknowledgments

Greatest appreciation should be given to the participant students of Animal Ecology Course (2006) for spending their time enthusiastically, with warm and hospitable thoughts during the field practice.

## References

- Beehler, B.M. and Mack, A.L. 1998. *Constraints to characterizing spatial heterogeneity in a lowland forest avifauna in New Guinea*. In: Adams, N.J. and Slotow, R.H. (Eds.). Proc. 22 Int Ornitholog. Congr. Durban: Birdlife South Africa. Pp. 2569–2579.
- Beehler, B.M., Pratt, T.K. and Zimmerman, D.A. 2001. *Burung-burung di Kawasan Papua* (Seri Panduan Lapangan LIPI). 497 pp.
- Dwiyanto, G. 1995. Jenis burung diurnal di Taman Wisata Alam Gunung Meja. *Skripsi Sarjana Kehutanan Fakultas Pertanian Universitas Cenderawasih Manokwari*.
- Leppe, D. and Tokede, M.J. 2006. *Potensi biofisik kawasan hutan Taman Wisata Alam Gunung Meja Manokwari*. Balai Penelitian Pengembangan Kehutanan Papua dan Maluku, Manokwari. 47 pp.
- Mayor, S. and Pattiselanno, F. 2008. Jenis burung di areal hutan mangrove Raja Ampat. *Warta Konservasi Lahan Basah*, Juli: 20–21.
- Pattiselanno, F. 2005. Birds diversity at the Mamberamo River Basin, Papua. *Biota*, X (3): 200–202.
- Rand, A.L. and Gilliard, E.T. 1967. *Handbook of New Guinea Birds*. The Trinity Press, Worcester, London. 612 pp.